

• •



2015 .

.....	2
.....	4
.....	4
.....	5
.....	5
.....	6
.....	7
.....	8
.....	9
.....	13
.....	13
.....	13

XVIII

» 14

2015 .

«

: . . „ - . -
 . „ „ . . .
 /
 , .
 , / :
 - .
 , / :
 , EMBASE MEDLINE.
 :
 -
 - ()
)
 :

1++	- , ()
1+	- , ,
1-	- , ,
2++	

	- .
2+	-
2-	-
3	(:
4	,)

, - ; :

, . :

:

	, 1++ , - , ,
	1+, , ,
	2++ , ,
1++ 1+	,
	2+, , , ;

[5,6,7,8].

Malassezia furfur

Malassezia [6,7].

[8].

[7].

- (« »)
-
- ()
-
-
-

-

, (crusta lactea –
).
 ,
 ; [7,11].
 :
 ,
 .
 ,
 [5,6,11].

- - .
 , , 3 .
 , ,
 [7].

- ()
 ,
 , .
 5 ,
 [6,7].
 6

[7,11].

1.

(8-12 .),

2.

3.

4.

1.

2.

3.

4.

5.

- / / (:
D07CA01), () [9,12]
2 5-7 .
(D) ,

- :
(D) 2 5
(D) 3-5 .

- / / (:
D07CA01) 2 (5-7
().

- (P06AB03) 3-5 3 :
()
- (P06AE07) 5 1 6
()

- :
-
-

, ,
,
:

- D07XC01) / (7-10
, 2- () Malassezia
, :
- 2% (J02AB02), 2
3-4 , 2
() [8,9,12,14]
, :
- 1-2% (D01AE13) 2
2-3 (D)
-1% , 0,2% - (D11AX12),
1 , 2 2-3
(D)
, :
- (D07AC13) 2
1 5-10 ,
()
- (D07AC01) 5-10
1 , 1-2
()
: / / (2
D07XC01) 2- , 2
5-10 ()
- / / (01)
, 2
5-10 ()
: 0,2% - (D11AX12)
1 , 2 10-14
(D)

- , 1-2 % (D01AE13) (D)

, , .

II,III

:

- (P06AE07) 10 1 7-10

()

- (R06AE09) 5 1 7-

10 ()

- (R06AX13) 10 1 7-10

()

- (R06AX27) 5 1 7-10 ()

,

:

- , . (J02AB02), 2 ,

15-30 100 1 , 30 -200

4 () [11,14]

- , . (J02AC02) 200 1

, 200 2 ()

- , . (D01BA02), 2 ,

40 250 1 4-6

()

- , . (J02AC01), 50 1

2 300 1 2 ()

1. I.Stefanaki, MD and A.Katsambas, MD. Therapeutic Update on Seborrheic Dermatitis. Skin therapy letter, 2010;15;5:1-7.
2. Ortonne J.P., Lacour J.P., Vitetta A., Le Fichoux Y. Comparative study of ketoconazole 2% foaming gel and betamethasone dipropionate 0.05% lotion in the treatment of seborrhoeic dermatitis in adults. Dermatology 1992; 184(4): 275-280.
3. Faergemann J. Seborrhoeic dermatitis and Pityrosporum orbiculare: treatment of seborrhoeic dermatitis of the scalp with miconazole-hydrocortisone (Daktacort), miconazole and hydrocortisone. Br J Dermatol. 1986; 114:695–700.
4. , 2013.
5. , 2013. 19, .186-189.
6. 20, .99-101.

7. . . . , . . .
1, 22, .235-242.
8.2009, .459-461.
9. Taieb A., Legrain V., Palmier C. et al. Topical ketoconazole for infantile seborrheic dermatitis. *Dermatologica*. 1990; 181:26–32.
Ford G.P., Farr P.M., Ive F.A., Shuster S. The response of seborrhoeic dermatitis to ketoconazole. *Br J Dermatol* 1984; 111: 603–607.
10. Prakash A., Benfield P. Topical mometasone. A review of its pharmacological properties and therapeutic use in the treatment of dermatological disorders. *Drugs* 1998; 55(1):145-163.
11. Naldi L, Rebora A. Clinical practice. Seborrheic dermatitis. *N Engl J Med* 360(4):387-96 (2009 Jan).
12. Crespo EV, Delgado FV. *Malassezia* species in skin disease. *Curr Opin Infect Dis* 15(2):133-42 (2002 Apr).
13. Meshkinpour A., Sun J., Weinstein G. An open pilot study using tacrolimus ointment in the treatment of seborrheic dermatitis. *J Am Acad Dermatol*. 2003; 49:145–147.
14. Wannanukul S., Chiabuncana J. Comparative study of 2% ketoconazole cream and 1% hydrocortisone cream in the treatment of infantile seborrheic dermatitis. *Med Assoc Thai* 2004; 87 (Suppl. 2):S68-S71,.
15. Metz M., Wahn U., Gieler U. et al. Chronic pruritus associated with dermatologic disease in infancy and childhood: update from an interdisciplinary group of dermatologists and pediatricians. *Pediatr Allergy Immunol*. 2013; 24(6):527–539.